



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

180. Proposed by the late JOSIAH H. DRUMMOND.

If r/s is such a value of p as makes $m/(p^2-2)$ integral, prove that $(3r+4s)/(2r+3s)$ is another such value, so that an indefinite number of integral values may be obtained.

Also, if r/s is such a value of p as makes $2m/(p^2-2)$ integral, prove that $2(r+s)/(r+2s)$ is also such a value.

GEOMETRY.

201. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Two plane sections of a right circular cone have their major axes AA' , aa' coplanar, and Aa on one generator equal to $A'a'$ on the other. The projections of the sections on any plane perpendicular to the axis are confocal.

202. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

The equations $1/(la) + 1/(m\beta) + 1/(n\gamma) = 0$ and $l\beta\gamma + m\alpha\gamma + n\beta\alpha = 0$ represent ellipses. If a, b, c are the sides of the triangle of reference, transform to Cartesian coördinates and find area of each ellipse.

CALCULUS.

166. Proposed by T. N. HAUN, Mohawk, Tenn.

Find the volume of the solid formed by the revolution of the curve $(y^2 + x^2) = a^2(x^2 - y^2)$ round the axis of x .

167. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Integrate,
$$\int_0^a \int_0^b \int_0^c \frac{z dx dy dz}{(x^2 + y^2 + z^2)^{\frac{3}{2}}}.$$

MECHANICS.

157. Proposed by T. W. WRIGHT, Schenectady, N. Y.

Explain why a waterfall h feet high can support a column of water $2h$ feet high.

158. Proposed by G. H. HARVILL, A. M., Malakoff, Texas.

Show that a law of density for points in space may be assumed such that the joint mass of any two points which are electrical images of each other in respect to a given sphere may be constant, and that their centers of gravity shall lie on the surface of the sphere.

AVERAGE AND PROBABILITY.

143. Proposed by L. C. WALKER, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

The extremities of two equal lines drawn from a fixed point in the circumference of a given circle is joined. Find the average area of the circle inscribed in the triangle formed.